

Question: Should BPD/DS vs. sleeve gastrectomy be used for weight loss?

№ of studies	Study design	Certainty assessment					№ of patients		Effect		Certainty	Importance
		Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	BPD/DS	sleeve gastrectomy	Relative (95% CI)	Absolute (95% CI)		
30-day morbidity												
2	observational studies	not serious	not serious	not serious	very serious ^a	none	10/267 (3.7%)	6/674 (0.9%)	OR 5.10 (0.24 to 107.62)	35 more per 1.000 (from 7 fewer to 483 more)	⊕○○○ VERY LOW	CRITICAL
30-day mortality												
2	observational studies	not serious	not serious	not serious	very serious ^a	none	0/267 (0.0%)	0/674 (0.0%)	not estimable	0 fewer per 1.000 (from 10 fewer to 10 more)	⊕○○○ VERY LOW	CRITICAL
Overall mortality												
1	observational studies	not serious	not serious	not serious	very serious ^b	none	0/100 (0.0%)	0/100 (0.0%)	RD 0.00 (-0.01 to 0.01)	0 fewer per 1.000 (from 20 fewer to 20 more)	⊕○○○ VERY LOW	CRITICAL

CI: Confidence interval; **OR:** Odds ratio